

Rates of Traumatic Injury by Idaho County — 2017–2021

August 2023

A Publication of the
Idaho Time-Sensitive Emergency Registry



Idaho TSE Registry
Trauma, Stroke, STEMI

Editors

Bozena M. Morawski, PhD, MPH, Epidemiologist
Randi K. Rycroft, MSPH, CTR, Registry Manager

Contributors

Regina Eck, Database Administrator
Michelle Barnett, CSTR, RHIT, CAISS, Idaho TSE Registry Project Specialist
Rosa Inskeep, CSTR, Idaho TSE Registry Trauma Registrar
Cheryl Hansen, CAISS, CSTR, Trauma Registrar
Louvier Gomez, CAISS, Trauma Registrar

Idaho Time-Sensitive Emergency Registry

P.O. Box 1278
Boise, Idaho 83701-1278
Phone: (208) 338-5100
Fax: (208) 344-0180
<https://idahotseregistry.org/>



Table of Contents

Preface and Acknowledgments.....	1
Preface	1
Acknowledgements.....	1
Background	2
Technical Notes.....	2
Participating Acute Care Facilities.....	2
Age-adjusted and Crude Incidence Rates	3
Confidence Intervals	3
Classification of Injury Mechanism and Intent.....	3
Limitations	3
I: Maps and Charts of Age-Adjusted Incidence by County and Mechanism of Injury, 2017–2021 Injury Years.....	4
Male and Female Idaho Residents	4
Male Idaho Residents.....	19
Female Idaho Residents.....	34
II: Maps and Charts of Crude Incidence by County and Mechanism of Injury, 2017–2021 Injury Years.....	49
Male and Female Idaho Residents	49
Male Idaho Residents.....	64
Female Idaho Residents	79
III: Maps and Charts of Age-Adjusted Incidence by County and Mechanism of Injury among Pediatric Patients, 2017–2021 Injury Years	94
Male and Female Pediatric (< 15 Years) Idaho Residents	94
Male Pediatric (< 15 Years) Idaho Residents.....	109
Female Pediatric (< 15 Years) Idaho Residents	124
IV: Maps and Charts of Age-Adjusted Incidence by County and Injury Intent, 2017–2021 Injury Years.....	139
Male and Female Idaho Residents	139
Male Idaho Residents.....	145
Female Idaho Residents	151
V: Maps and Charts of Crude Incidence by County and Injury Mechanism by Location of Injury, 2017–2021 Injury Years	157
Male and Female Patients (Idaho and non-Idaho Residents)	157

Male Patients (Idaho and non-Idaho Residents)	172
Female Patients (Idaho and non-Idaho Residents)	187
References	202
Appendices	203
Appendix A: Map of Idaho TSE Districts and Counties	203
Appendix B: 2000 United States Standard Population.....	204
Appendix C: 2017–2021 State of Idaho Population by County	205

Preface and Acknowledgments

Preface

The Time-Sensitive Emergency (TSE) Registry was established by Idaho Code §57-2001 to collect the data required to analyze the incidence, severity, causes, costs, and outcomes of traumatic injury, stroke, and ST-elevation myocardial infarction (STEMI) in Idaho. TSE Registry trauma data describe the nature and scope of traumatic injury, severity of injury, performance of out-of-hospital and hospital emergency medical systems, and patient outcomes for Idaho residents and non-Idaho residents treated for a traumatic injury at an Idaho acute care facility.

“Rates of Traumatic Injury by Idaho County — 2017–2021” shows, via charts and maps, where Idaho resident traumatic injury patients reside at the time of their injury and where in Idaho traumatic injuries occur among Idaho residents and non-residents. This document includes data from traumatic events occurring during 2017–2021.

We hope that these data are used by the Idaho Time Sensitive Emergency (TSE) Council, public health officials, hospital administrators, physicians, and others to develop prevention and intervention strategies, and identify high-risk populations in Idaho within the constraints of sex- and county-specific data.

Acknowledgements

The Idaho Department of Health and Welfare, Bureau of Emergency Medical Services and Preparedness (BEMSP) contracts with, and provides funding to, the Idaho Hospital Association (IHA) to maintain a statewide trauma registry.

The statewide TSE Registry database is a product of collaboration among multiple report sources, including Idaho’s acute care hospitals. Their reporting of timely, accurate, and complete data describing traumatic injury is acknowledged and sincerely appreciated.

The TSE Registry also thanks the Division of Public Health, Idaho Department of Health and Welfare for its continued partnership and collaboration.

SUGGESTED CITATION: Morawski BM, Rycroft RK. *Rates of Traumatic Injury by Idaho County — 2017–2021*. Boise, ID: Time-Sensitive Emergency Registry; August 2023.

COPYRIGHT INFORMATION: All material in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Background

During 2017–2021, 38,592 qualifying traumatic injuries were reported to the Idaho TSE Registry – a 9.6% increase over the 2016–2020 injury years. Incident rates in this document were calculated using two types of numerators:

1. attributing events to a specific Idaho county using patient ZIP code of residence at time of injury; and
2. attributing events to a specific Idaho county using injury location ZIP code.

Patient ZIP code of residence at time of injury was missing or the patient was not a current resident of the United States in 10 (0.03%) of qualifying traumatic injuries. Of the 38,592 qualifying traumatic injuries, 32,037 occurred among Idaho residents and are depicted in the maps herein describing traumatic injury by patient residency.

Injury location ZIP code was missing (N = 32) or occurred out of state (N = 4,824) for 4,856 events (12.6%) occurring during 2017–2021. Most injuries with reported location occurred in Idaho (87% or 33,736 events). Also treated in Idaho acute care facilities were injuries occurring in Oregon (2.2% or 836 events), Wyoming (1.4% or 537 events), Washington (0.9% or 358 events), Montana (0.5% or 182 events), Nevada (0.3% or 99 events) and Utah (0.1% or 29 events).

Technical Notes

Participating Acute Care Facilities

As of December 31, 2020, 42 of Idaho's acute care facilities were reporting trauma cases to TSE Registry. We thank these facilities for their participation.

- Bear Lake Memorial Hospital
- Benewah Community Hospital
- Bingham Memorial Hospital
- Bonner General Health
- Boundary Community Hospital
- Caribou Memorial Hospital, Living Center & Clinics
- Cascade Medical Center
- Cassia Regional Medical Center
- Clearwater Valley Hospital & Clinics
- Eastern Idaho Regional Medical Center
- Franklin County Medical Center
- Gritman Medical Center
- Idaho Falls Community Hospital
- Kootenai Health
- Lost Rivers Medical Center
- Madison Memorial Hospital
- Minidoka Memorial Hospital
- Nell J. Redfield Memorial Hospital
- North Canyon Medical Center
- Portneuf Medical Center
- Power County Hospital District
- Saint Alphonsus Regional Medical Center
- Saint Alphonsus Medical Center – Nampa
- Shoshone Medical Center
- St. Joseph Regional Medical Center
- St. Luke's Boise Medical Center
- St. Luke's Children's
- St. Luke's Elmore
- St Luke's Fruitland
- St. Luke's Jerome
- St. Luke's Magic Valley Regional Medical Center
- St. Luke's McCall
- St Luke's Meridian Medical Center
- St. Luke's Nampa Medical Center
- St. Luke's Wood River Medical Center
- St. Mary's Hospital & Clinics
- Steele Memorial Medical Center
- Syringa Hospital & Clinics
- Teton Valley Health Care, Inc.
- Valor Health
- Weiser Memorial Hospital
- West Valley Medical Center

Age-adjusted and Crude Incidence Rates

Age-adjusted incidence rates shown on the maps included in this document were calculated using the direct method of standardization, and standardized to the age distribution of the 2000 U.S. population (see Appendix B). Incidence rates represent the average number of traumatic injuries annually per 100,000 persons. Age adjustment allows rates from one geographic area or time period to be compared with rates from other geographic areas or time periods that may have differences in age distributions. Any observed differences in age-adjusted incidence rates between populations are not due to differing age structures.

Crude incidence rates shown on the maps included in this document were calculated as the number of traumatic injuries occurring among residents of that county (or the number of injuries occurring in that county) and divided by the population of that county for the injury year. No standardization, or age-adjustment, was applied to these rates. Crude rates are useful for public health by identifying areas of higher absolute burden of injury-related morbidity and mortality, scaled to population.

Rates are sorted and displayed by color groups. Counties are assigned to one of five color groups using a clustering method based on Euclidean distances, whereby the center of each cluster is based on least squares estimation. This approach is commonly referred to as a *k*-means model.¹

Confidence Intervals

Confidence intervals, which are estimated from available data, provide a range of values that are likely to include the true and unknown population value. The width of a confidence interval is a measure of variability, with wider confidence intervals connoting less reliable estimates. The charts included in this document display confidence intervals around county-specific rate estimates.

Classification of Injury Mechanism and Intent

Maps in this document are presented by injury mechanism and intent. Injury mechanism is typically thought of as the “how” of injury, e.g., firearm, motor vehicle accident, fall. Injury intent can be thought of as the manner in which a person was injured or died, i.e., unintentionally, intentionally (homicide/assault, suicide/self-harm) or undetermined. Injury mechanism and injury intent as shown in this document are classified based on the ICD-10-CM external cause of injury code. For non-fatal injuries, external cause codes are classified as described in the “Proposed framework for presenting injury data using the International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM) diagnosis codes.”² Fatal injuries are classified as described in the “The ICD-10 injury mortality diagnosis matrix”^{3,4} based on ICD-10 mortality codes.


Limitations

Maps in this document based on injury location use numerator data that include Idaho residents and non-residents in conjunction with denominator data that only represent Idaho residents. Thus, the rates for a given county may be artificially high (in particular when comparing these data to other counties) in areas where there are large numbers of non-Idaho residents experiencing traumatic injury, e.g. state border communities and counties, counties with relatively large number of tourists. These rates may be particularly inflated in areas with relatively small resident populations. For all of these reasons, rates of injury based on injury location should be interpreted with prudence.


These data are, however, being presented as rates of injury based on location as they provide some insight as to the burden of injury that first responders and local hospitals may experience with respect to treating traumatic injury. They may also be hypothesis generating with respect to public health intervention in a particular county.

I: Maps and Charts of Age-Adjusted Incidence by County and Mechanism of Injury, 2017–2021 Injury Years


Male and Female Idaho Residents




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
All causes (known and unknown)
Males and Females
State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Total with known cause
Males and Females
State of Idaho, by County, 2017-2021




*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Bites and Stings
Males and Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 9.3




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Cut/pierce

Males and Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 2.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Drowning/submersion
Males and Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Fall


Males and Females

State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 4.8




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Fire/burn

Males and Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 20.9




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Firearm
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 3.9




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Machinery
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 65.4




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Motor vehicle traffic
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 3.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Natural/environmental
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 6.4




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Other Specified
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 5.5




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Suffocation
Males and Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 2.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Unspecified

Males and Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

Male Idaho Residents




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
All causes (known and unknown)
Males
State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 418.3




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Total with known cause

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Bites and Stings
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 13.5




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Cut/pierce

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.5




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Drowning/submersion
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Fall

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 6.5




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Fire/burn

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 35.1




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Firearm
Males
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 7.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Machinery
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 83.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Motor vehicle traffic

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 4.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Natural/environmental
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 9.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Other Specified

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 8.2




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Suffocation
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 2.4




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Unspecified

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*


Female Idaho Residents




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
All causes (known and unknown)
Females
State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Total with known cause
Females
State of Idaho, by County, 2017-2021




*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Bites and Stings
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 5.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Cut/pierce
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Drowning/submersion
Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
 Arrows indicate upper confidence interval extends beyond graphic area.




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Fall

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.2




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Fire/burn

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 6.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Firearm

Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 0.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Machinery
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 47.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Motor vehicle traffic
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.5




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Natural/environmental
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
 Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Other Specified
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.9




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Suffocation
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
 Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.6



Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Unspecified
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.


*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.*

II: Maps and Charts of Crude Incidence by County and Mechanism of Injury, 2017–2021 Injury Years

Male and Female Idaho Residents




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
All causes (known and unknown)
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Total with known cause
Males and Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.6




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Bites and Stings

Males and Females

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.


Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 9.0




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Cut/pierce
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 2.5




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Drowning/submersion
Males and Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 197.8




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Fall

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Fire/burn

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 20.7




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Firearm

Males and Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Machinery
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 65.8




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Motor vehicle traffic
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 3.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Natural/environmental
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 6.3




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Other Specified
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 5.3




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Suffocation
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.0




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Unspecified
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.*


Male Idaho Residents




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
All causes (known and unknown)
Males
State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
 Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Total with known cause
Males
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.6




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Bites and Stings
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.*

State of Idaho Rate: 13.2




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Cut/pierce

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.5




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Drowning/submersion
Males
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Fall

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 6.6




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Fire/burn

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 34.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Firearm

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 7.0




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Machinery

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 83.8




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Motor vehicle traffic

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Natural/environmental

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 9.7




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Other Specified

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 7.7




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury


Suffocation

Males

State of Idaho, by County, 2017-2021



State of Idaho Rate: 2.3




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Unspecified

Males

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

Female Idaho Residents

State of Idaho Rate: 325.2




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

All causes (known and unknown)


Females

State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Total with known cause

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 1.7




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Bites and Stings

Females

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.

Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 4.8




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Cut/pierce

Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.5




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Drowning/submersion
Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.*

State of Idaho Rate: 221.2




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Fall

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.2




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Fire/burn

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 6.5




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Firearm

Females

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.


Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 0.7




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury
Machinery
Females
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 47.6




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Motor vehicle traffic

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 2.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Natural/environmental

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.9




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Other Specified

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.8




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Suffocation

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.7




Crude Incidence Rates - Calculated from Patient Residence at Time of Injury

Unspecified

Females

State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.


Rates are not adjusted to a standard population. U.S. FIPS codes label counties.

Colors denote rate clustering. Width of bars indicate 95% confidence intervals.


Arrows indicate upper confidence interval extends beyond graphic area.

III: Maps and Charts of Age-Adjusted Incidence by County and Mechanism of Injury among Pediatric Patients, 2017–2021 Injury Years

Male and Female Pediatric (< 15 Years) Idaho Residents




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)
All causes (known and unknown)
Males and Females (< 15 years)
State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 131.4




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Total with known cause


Males and Females (< 15 years)

State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Bites and Stings

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Cut/pierce

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.7




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Drowning/submersion

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 53.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Fall

Males and Females (< 15 years)

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.

Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.9




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Fire/burn

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.1




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Firearm

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.0




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Machinery

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 15.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Motor vehicle traffic

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.7




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Natural/environmental

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Other Specified

Males and Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.1




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)


Suffocation

Males and Females (< 15 years)

State of Idaho, by County, 2017-2021



State of Idaho Rate: 1.3




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Unspecified

Males and Females (< 15 years)

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

Male Pediatric (< 15 Years) Idaho Residents

State of Idaho Rate: 162.6




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

All causes (known and unknown)

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 160.9




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Total with known cause

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.5




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Bites and Stings

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.3




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Cut/pierce

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 4.4




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Drowning/submersion

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 63.4




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Fall

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 5.5




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Fire/burn

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Firearm

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 1.6




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Machinery

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 17.2




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Motor vehicle traffic

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 2.1




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Natural/environmental

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.7




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Other Specified

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.7




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Suffocation

Males (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 1.7




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Unspecified

Males (< 15 years)

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

Female Pediatric (< 15 Years) Idaho Residents

State of Idaho Rate: 101.5




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

All causes (known and unknown)

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 100.5




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Total with known cause

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.5




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Bites and Stings

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.2




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Cut/pierce

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.9




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Drowning/submersion

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 43.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Fall

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.2




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Fire/burn

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.4




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Firearm

Females (< 15 years)


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 0.3




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)
Machinery
Females (< 15 years)
State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 14.3




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Motor vehicle traffic

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 1.2




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Natural/environmental

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.8




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Other Specified

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.5




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Suffocation

Females (< 15 years)


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 0.9




Age-Adjusted Incidence of Pediatric Injury (Idaho Residents)

Unspecified

Females (< 15 years)

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.


**IV: Maps and Charts of Age-Adjusted Incidence by County and Injury Intent, 2017–2021
Injury Years**

Male and Female Idaho Residents

State of Idaho Rate: 11.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Assault
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 0.8




**Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Other/Legal/War
Males and Females
State of Idaho, by County, 2017-2021**




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 24.7




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Self-harm/Suicide
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 318.8




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Unintentional
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.1



Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Undetermined
Males and Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

Male Idaho Residents

State of Idaho Rate: 16.5




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Assault

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Other/Legal/War
Males
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 39.0




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Self-harm/Suicide
Males
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Unintentional
Males
State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.6




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Undetermined

Males

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

Female Idaho Residents

State of Idaho Rate: 5.3




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Assault

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 0.1




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury

Other/Legal/War

Females


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 10.4




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Self-harm/Suicide
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

*Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Unintentional
Females
State of Idaho, by County, 2017-2021




*Idaho Time-Sensitive Emergency Registry.
 Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
 Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 0.5



Age-Adjusted Incidence - Calculated from Patient Residence at Time of Injury
Undetermined
Females
State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates age-adjusted to the 2000 U.S. population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

V: Maps and Charts of Crude Incidence by County and Injury Mechanism by Location of Injury, 2017–2021 Injury Years

Male and Female Patients (Idaho and non-Idaho Residents)

State of Idaho Rate: 356.0




Crude Incidence Rates - Calculated from Injury ZIP Code

All causes (known and unknown)

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 354.9




Crude Incidence Rates - Calculated from Injury ZIP Code

Total with known cause

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 1.3




Crude Incidence Rates - Calculated from Injury ZIP Code

Bites and Stings

Males and Females



State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.



State of Idaho Rate: 7.7




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.7






Crude Incidence Rates - Calculated from Injury ZIP Code

Fall

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 4.6




Crude Incidence Rates - Calculated from Injury ZIP Code

Fire/burn

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 20.3




Crude Incidence Rates - Calculated from Injury ZIP Code

Firearm

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 3.3




Crude Incidence Rates - Calculated from Injury ZIP Code

Machinery

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 66.2




Crude Incidence Rates - Calculated from Injury ZIP Code

Motor vehicle traffic

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.5




Crude Incidence Rates - Calculated from Injury ZIP Code

Natural/environmental

Males and Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 5.7




Crude Incidence Rates - Calculated from Injury ZIP Code

Other Specified

Males and Females



State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.


State of Idaho Rate: 5.1



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 1.2




Crude Incidence Rates - Calculated from Injury ZIP Code

Unspecified

Males and Females

State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

Male Patients (Idaho and non-Idaho Residents)

State of Idaho Rate: 399.6




Crude Incidence Rates - Calculated from Injury ZIP Codes

All causes (known and unknown)

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 398.1




Crude Incidence Rates - Calculated from Injury ZIP Code

Total with known cause

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.


Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 1.3





Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.





State of Idaho Rate: 11.3



Idaho Time-Sensitive Emergency Registry.


Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 3.8



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.




Crude Incidence Rates - Calculated from Injury ZIP Code

Fall

Males



State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.


State of Idaho Rate: 6.2



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 34.0




Crude Incidence Rates - Calculated from Injury ZIP Code

Firearm

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 6.1




Crude Incidence Rates - Calculated from Injury ZIP Code

Machinery

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 84.2




Crude Incidence Rates - Calculated from Injury ZIP Code

Motor vehicle traffic

Males



State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.


State of Idaho Rate: 4.4



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 8.5




Crude Incidence Rates - Calculated from Injury ZIP Code

Other Specified

Males


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.


Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 7.4




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.



State of Idaho Rate: 1.5




Crude Incidence Rates - Calculated from Injury ZIP Code

Unspecified

Males


State of Idaho, by County, 2017-2021




Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.


Female Patients (Idaho and non-Idaho Residents)




**Crude Incidence Rates - Calculated from Injury ZIP Code
All causes (known and unknown)
Females
State of Idaho, by County, 2017-2021**



*Idaho Time-Sensitive Emergency Registry.
Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*




Crude Incidence Rates - Calculated from Injury ZIP Code Total with known cause Females State of Idaho, by County, 2017-2021



*Idaho Time-Sensitive Emergency Registry.
Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.*

State of Idaho Rate: 1.4




Crude Incidence Rates - Calculated from Injury ZIP Code

Bites and Stings

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 4.1




Crude Incidence Rates - Calculated from Injury ZIP Code

Cut/pierce

Females



State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.


State of Idaho Rate: 1.5



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 212.1




Crude Incidence Rates - Calculated from Injury ZIP Code

Fall

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals.

State of Idaho Rate: 2.9




Crude Incidence Rates - Calculated from Injury ZIP Code

Fire/burn

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 6.5




Crude Incidence Rates - Calculated from Injury ZIP Code

Firearm

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 0.6




Crude Incidence Rates - Calculated from Injury ZIP Code

Machinery

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.


Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 48.1





Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.




State of Idaho Rate: 2.6



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties.
Colors denote rate clustering. Width of bars indicate 95% confidence intervals.
Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.8




Crude Incidence Rates - Calculated from Injury ZIP Code

Other Specified

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 2.7




Crude Incidence Rates - Calculated from Injury ZIP Code

Suffocation

Females


State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

State of Idaho Rate: 0.8




Crude Incidence Rates - Calculated from Injury ZIP Code

Unspecified

Females

State of Idaho, by County, 2017-2021



Idaho Time-Sensitive Emergency Registry.

Rates are not adjusted to a standard population. U.S. FIPS codes label counties. Colors denote rate clustering. Width of bars indicate 95% confidence intervals. Arrows indicate upper confidence interval extends beyond graphic area.

References

1. SAS Institute Inc. SAS/STAT® 13.2 User's Guide. In: Cary, NC: SAS Institute Inc.; 2014.
2. Hedegaard H, Johnson RL, Warner M, Chen H, Annest JL. Proposed Framework for Presenting Injury Data Using the International Classification of Diseases, Tenth Revision, Clinical Modification (ICD–10–CM) Diagnosis Codes. *National Health Statistics Reports*. 2016(89).
3. Fingerhut LA, Warner M. The ICD-10 injury mortality diagnosis matrix. *Inj Prev*. 2006;12(1):24-29.
4. Corrections. *Injury Prevention*. 2006;12(2):135-135.

Appendices

Appendix A: Map of Idaho TSE Districts and Counties



Courtesy of Idaho Department of Health and Welfare, accessible from:

<https://idaho.maps.arcgis.com/apps/View/index.html?appid=33800b0a1cdf4e748b20dbbdacf2dbbc>

Appendix B: 2000 United States Standard Population

Age Group	2000 US Standard Population (Census P25-1130)
0	3,794,901
1-4	15,191,619
5-9	19,919,840
10-14	20,056,779
15-19	19,819,518
20-24	18,257,225
25-29	17,722,067
30-34	19,511,370
35-39	22,179,956
40-44	22,479,229
45-49	19,805,793
50-54	17,224,359
55-59	13,307,234
60-64	10,654,272
65-69	9,409,940
70-74	8,725,574
75-79	7,414,559
80-84	4,900,234
85+	4,259,173
Total	274,633,642

Source: SEER Program, National Cancer Institute, 2021.

Appendix C: 2017–2021 State of Idaho Population by County

Geographic Area	2017	2018	2019	2020	2021	Total
State of Idaho	1,719,745	1,752,074	1,789,060	1,826,913	1,900,923	8,988,715
Ada	456,994	469,161	481,880	494,399	511,931	2,414,365
Adams	4,128	4,215	4,265	4,447	4,625	21,680
Bannock	85,571	86,761	87,822	88,795	88,263	437,212
Bear Lake	6,029	6,026	6,131	6,143	6,545	30,874
Benewah	9,167	9,222	9,303	9,430	9,931	47,053
Bingham	45,888	46,054	46,772	47,202	48,876	234,792
Blaine	22,401	22,736	23,012	23,426	24,766	116,341
Boise	7,370	7,689	7,903	8,065	8,094	39,121
Bonner	43,715	44,709	45,774	46,817	49,491	230,506
Bonneville	114,608	116,661	119,286	122,134	127,930	600,619
Boundary	11,995	12,073	12,340	12,656	12,561	61,625
Butte	2,582	2,592	2,620	2,646	2,654	13,094
Camas	1,092	1,098	1,091	1,130	1,139	5,550
Canyon	217,037	223,518	230,270	237,053	243,115	1,150,993
Caribou	6,980	7,019	7,149	7,123	7,111	35,382
Cassia	23,661	23,769	24,052	24,277	25,164	120,923
Clark	878	845	838	852	792	4,205
Clearwater	8,651	8,759	8,761	8,846	8,895	43,912
Custer	4,138	4,216	4,273	4,249	4,428	21,304
Elmore	26,907	27,309	27,373	27,448	28,827	137,864
Franklin	13,501	13,718	13,898	14,215	14,666	69,998
Fremont	13,140	13,162	13,105	13,218	13,592	66,217
Gem	17,351	17,622	18,174	18,703	19,792	91,642
Gooding	15,155	15,187	15,290	15,618	15,772	77,022
Idaho	16,392	16,469	16,623	16,823	17,040	83,347
Jefferson	28,448	29,418	29,918	30,581	32,202	150,567
Jerome	23,791	24,078	24,486	24,578	24,662	121,595
Kootenai	157,594	161,279	165,656	170,628	179,789	834,946
Latah	39,903	39,996	40,372	40,830	40,313	201,414
Lemhi	7,837	7,974	8,067	8,054	8,162	40,094
Lewis	3,883	3,835	3,845	3,838	3,715	19,116
Lincoln	5,361	5,360	5,325	5,358	5,282	26,686
Madison	39,540	39,451	40,205	40,318	53,881	213,395
Minidoka	20,683	20,667	20,910	21,216	21,955	105,431
Nez Perce	40,376	40,408	40,601	40,755	42,454	204,594
Oneida	4,393	4,429	4,500	4,520	4,611	22,453
Owyhee	11,632	11,664	11,823	12,133	12,336	59,588
Payette	23,218	23,560	24,051	24,771	26,350	121,950
Power	7,587	7,654	7,636	7,643	7,950	38,470
Shoshone	12,511	12,781	12,868	12,911	13,612	64,683
Teton	11,463	11,640	12,096	12,501	12,267	59,967
Twin Falls	85,439	86,166	87,127	88,411	92,243	439,386
Valley	10,700	11,054	11,443	11,792	12,241	57,230
Washington	10,055	10,070	10,126	10,360	10,898	51,509